



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

365. Proposed by C. N. SCHMALL, New York City.

Show that the area inclosed by each of the following three curves is equal to the circle of radius a ; viz., πa^2 .

$$(1) \quad a^2 x^2 = y^2(2a - y), \quad (2) \quad a^2 - x^2 = (y - mx^2)^2, \quad (3) \quad (xy + c + bx^2)^2 = x^2(a^2 - x^2).$$

I. SOLUTION BY A. M. HARDING, University of Arkansas.

If we change these equations to parametric forms we obtain

$$(1) \quad x = 4a \cos^3 t \sin t, \quad y = 2a \cos^2 t,$$

$$(2) \quad x = a \sin t, \quad y = ma^2 \sin^2 t + a \cos t,$$

$$(3) \quad x = a \sin t, \quad y = \frac{a^2 \cos t - a^2 b \sin t - c \csc t}{a}.$$

Hence,

$$(1) \quad \text{Area} = \int y \, dx = \int_0^\pi 8a^2(4 \cos^6 t - 3 \cos^4 t) dt = \pi a^2;$$

$$(2) \quad \text{Area} = \int y \, dx = \int_0^{2\pi} (ma^2 \sin^2 t + a \cos t) a \cos t \, dt = \pi a^2; \text{ and}$$

$$(3) \quad \text{Area} = \int y \, dx = \int_0^{2\pi} (a^2 \cos t - a^2 b \sin t - c \csc t) \cos t \, dt = \pi a^2.$$

II. SOLUTION BY H. C. FEEMSTER, York College, Nebraska.

$$A_1 = \int_0^{2a} \int_{-(y/a)\sqrt{2ay-y^2}}^{(y/a)\sqrt{2ay-y^2}} dy \, dx = \left[-\frac{3a^2 + ax - 2x^2}{3} \sqrt{2ax - x^2} + a^2 \operatorname{vers}^{-1} \frac{x}{a} \right]_0^{2a} = \pi a^2;$$

$$A_2 = \int_{-a}^a \int_{mx - \sqrt{a^2 - x^2}}^{mx + \sqrt{a^2 - x^2}} dx \, dy = \left[x \sqrt{a^2 - x^2} + a^2 \sin^{-1} \frac{x}{a} \right]_{-a}^a = \pi a^2;$$

and

$$A_3 = \int_{-a}^a \int_{-\sqrt{a^2 - x^2} - (c/x) - bx}^{\sqrt{a^2 - x^2} - (c/x) - bx} dx \, dy = \left[x \sqrt{a^2 - x^2} + a^2 \sin^{-1} \frac{x}{a} \right]_{-a}^a = \pi a^2.$$

NUMBER THEORY.**218. Proposed by ELIJAH SWIFT, University of Vermont.**

If p is prime > 3 show that

$$\sum_{a=1}^{a=p-1} \frac{1}{a^2} \equiv 0 \pmod{p}. \quad (1)$$

I. SOLUTION BY TRACY A. PIERCE, Berkeley, Cal.

In (1), we may replace 1 by a^{p-1} , since $a^{p-1} \equiv 1 \pmod{p}$. We then have

$$\sum_{a=1}^{a=p-1} a^{p-3} \equiv 0 \pmod{p}.$$

But it is well known that the sum of like powers of the numbers 1, 2, 3, ..., $p - 1$ is divisible by p if the power is not a multiple of $p - 1$; hence, the theorem is proved.

As a generalization of the congruence above, we may state

$$\sum_{a=1}^{a=p-1} \frac{1}{a^k} \equiv 0 \pmod{p}$$

if k is not a multiple of $p - 1$.

II. SOLUTION BY THE PROPOSER.

$$\sum_{a=1}^{a=p-1} \frac{1}{a^2} \equiv \sum_{a=1}^{a=p-1} a^2 \pmod{p}.$$

But

$$\sum_{a=1}^{a=p-1} a^2 \equiv A_1^2 - 2A_2$$

where

$$A_1 = \sum_{a=1}^{p-1} a, \quad A_2 = \sum_{\substack{a_1=1 \\ a_2=1}}^{p-1} a_1 a_2, \quad a_1 \neq a_2.$$

In Bachmann's *Niedere Zahlentheorie*, Vol. I, page 155, it is proved that A_1 and A_2 are divisible by p , whence the theorem.

NOTES AND NEWS.

EDITED BY W. D. CAIRNS.

Professor ANDREW W. PHILLIPS, of Yale University, died on January 20, 1915. He was joint author of Phillips and Fisher's *Geometry*.

Professor R. M. BARTON of the University of New Mexico has been appointed professor of mathematics in Lombard College.

Professor W. H. ROEVER, representative of Washington University on the editorial staff of the MONTHLY, was elected to honorary membership in the Society of Phi Beta Kappa by the Washington University Chapter, which was installed last year.

"The arithmetic mean as approximately the most probable value *a posteriori* under the gaussian probability law," by EDWARD L. DODD, is the title of a pamphlet published by the University of Texas as its January, 1915, *Bulletin*.

The February number of the *Proceedings of the National Academy of Sciences* contains two mathematical articles. The first is by Professor E. J. WILCZYNSKI and bears the title "Conjugate systems of space curves." The second is due to